arXiv:hep-th/0606151v5 10 Jul 2008

Frobenius transformation, mirror map and instanton numbers

Albert Schwarz, Vadim Vologodsky

July 10, 2008

Abstract

We show that one can express Frobenius transformation on middle-dimensional p-adic cohomology of Calabi-Yau threefold in terms of mirror map and instanton numbers. We express the mirror map in terms of Frobenius transformation on p-adic cohomology . We discuss a p-adic interpretation of the conjecture about integrality of Gopakumar-Vafa invariants.

1 Introduction

In [1] we gave an expression of instanton numbers in terms of Frobenius transformation on middle-dimensional p-adic cohomology of Calabi-Yau threefold with respect to canonical coordinates on moduli space of complex structures . We mentioned that conversely Frobenius map in canonical coordinates can be expressed in terms of instanton numbers. (A rigorous exposition of the results of [1] was given in [2].) In present paper we will prove that Frobenius map on middle-dimensional cohomology constructed by means of any coordinate system in the neighborhood of maximally unipotent boundary point of complex structures can be expressed in terms of mirror map and instanton numbers (Sec.2) The calculations in Sec. 2 generalize some calculations of [1], however, our exposition is independent of [1]. In Sec. 3 we will discuss integrality of mirror map ; we will give an explicit formula for mirror map in terms of Frobenius transformation on cohomology of Calabi-Yau manifold. (Talking about Frobenius map on cohomology we have in mind the middle-dimensional cohomology with coefficients in the ring $\mathbb{Z}_p$ of integer p-adic numbers; see [3] for transparent construction of this map.) In Sec.4 we interpret the conjecture about integrality of Gopakumar-Vafa invariants in terms of Frobenius map on p-adic wave functions. We hope that this interpretation will allow us to prove the conjecture. In the last section we describe a way that permits us to apply our calculation of Frobenius map to computation of zeta-functions of Calabi-Yau manifolds over finite fields. These zeta-functions were studied in [8]; our calculations are more general , but less explicit.

2 Calculation of Frobenius map

Let us consider a family of complex Calabi-Yau manifolds of dimension n in the neighborhood of maximally unipotent boundary point in the moduli space $\mathcal{M}$ of complex structures. The coordinates in the neighborhood of the boundary point will be denoted by $z_1, ..., z_r$; the points on the boundary divisor obey $z_i = 0$. We assume that the moduli space $\mathcal{M}$ is defined over $\mathbb{Z}$.

The corresponding B-model can be formulated in terms of Hodge filtration $\mathcal{F}^p$, monodromy weight filtration $\mathcal{W}_k$ and Gauss-Manin connection ∇ on the bundle of cohomology groups H^n (with complex coefficients). (See, for example, Sec 5.1 of [4] .) We assume that Morrrison integrality conjecture (see [5] or Sec 5.2.2 of [4]) is satisfied.

We will consider Calabi -Yau threefolds. In this case the cohomology group H^3 can be represented as a direct sum of the subgroups $I^{k,k} = \mathcal{W}_{2k} \bigcap \mathcal{F}^k$ where $k = 0, 1, 2, 3$, and $\text{rank} I^{0,0} = \text{rank} I^{3,3} = 1, \text{rank} I^{1,1} = \text{rank} I^{2,2} = r$. (Here r stands for the dimension of $\mathcal{M}$.)

Let us assume for simplicity that $r = 1$. Then we can take a symplectic basis of H^3 consisting of vectors $e_k \in I^{3-k,3-k}$, $, k = 0, 1, 2, 3$. (In other words, the vectors e_k obey $< e_3, e_0 >= 1, < e_2, e_1 >= -1$.)

For appropriate choice of this basis (see, for example [4]) Gauss- Manin connection takes the form

$$\nabla_\delta e_3 = 0, \tag{1}$$

$$\nabla_\delta e_2 = Y_3(z) e_3, \tag{2}$$

$$\nabla_\delta e_1 = Y_2(z) e_2, \tag{3}$$

$$\nabla_\delta e_0 = Y_1(z) e_1. \tag{4}$$

Here δ stands for logarithmic derivative with respect to coordinate z in the neighborhood of maximally unipotent boundary point. Compatibility of Gauss-Manin connection with symplectic structure on cohomology implies that $Y_1 = Y_3$.

We follow the notations of Sec 5.6.3 of [4]. It is shown in this section that the mirror map $q(z)$ can be expressed in terms of $Y_1 = Y_3$; namely $\delta \log q = Y_3$. In canonical coordinate q we have $Y_1 = Y_3 = 1$ (for appropriate normalization of the basis in cohomology) and Y_2 can be expressed in terms of instanton numbers (up to a constant summand.) In general Y_i can be expressed in terms of mirror map and instanton numbers. There exists also an expression of Y_i in terms of solutions of Picard-Fuchs equations (of periods of holomorphic 3-form $\Omega = f e_0$.) We will need the relation

$$\delta \log(f^4 Y_1^3 Y_2^2 Y_3) = c_3, \tag{5}$$

where c_3 stands for one of the coefficients of Picard-Fuchs equation

$$\nabla_\delta^4 \Omega = c_3 \nabla_\delta^3 \Omega + c_2 \nabla_\delta^2 \Omega + c_1 \nabla_\delta \Omega + c_0 \Omega. \tag{6}$$

To obtain (5) we substitute $\Omega = fe_0$ into Picard-Fuchs equation and apply (1)-(4). We will use (5) to calculate $f^2Y_1^2Y_2$.

One can prove that the action of Gauss-Manin connection on cohomology with coefficients in $\mathbb{Z}_p$ is given by (1)-(4) where Y_i are considered as series with p-adic coefficients ;see [2]. (In this statement one should assume that the canonical coordinate and the vectors e_i are normalized in appropriate way.)

The map $z \to z^p$ is called Frobenius map; it induces a map on functions of z denoted by Fr. It is a non-trivial fact that it can be lifted to a map of cohomology groups with coefficients in $\mathbb{Z}_p$; this map (Frobenius transformation) also will be denoted by Fr. Frobenius transformation is compatible with symplectic structure on cohomology:

$$< \mathrm{Fr}a, \mathrm{Fr}b >= p^3 \mathrm{Fr} < a, b >, \tag{7}$$

where $< a, b >$ stands for the inner product of cohomology classes.

The Frobenius transformation does not preserve the Hodge filtration; however, for $p > 3$ it satisfies

$$\mathrm{Fr}\mathcal{F}^s \subset p^s \mathcal{F}^0. \tag{8}$$

We will always impose the condition $p > 3$.

The Frobenius map preserves the monodromy weight filtration; hence its matrix is triangular in our basis:

$$\mathrm{Fr}e_3 = a_{33}e_3, \tag{9}$$

$$\mathrm{Fr}e_2 = pa_{22}e_2 + pa_{32}e_3, \tag{10}$$

$$\mathrm{Fr}e_1 = p^2a_{11}e_1 + p^2a_{21}e_2 + p^2a_{31}e_3, \tag{11}$$

$$\mathrm{Fr}e_0 = p^3a_{00}e_0 + p^3a_{10}e_1 + p^3a_{20}e_2 + p^3a_{30}e_3. \tag{12}$$

Notice that in the LHS of these formulas the vectors e_i are calculated at the point z and in the RHS these vectors are calculated at the point z^p. The coefficients belong to $\mathbb{Z}_p[[z]]$.

From the relation (7) we obtain

$$a_{33}a_{00} = 1, a_{22}a_{11} = 1, a_{00}a_{32} = a_{22}a_{10}, \tag{13}$$

$$a_{11}a_{20} - a_{21}a_{10} + a_{31}a_{00} = 0 \tag{14}$$

The Frobenius map is compatible with Gauss-Manin connection:

$$\nabla_\delta \mathrm{Fr} = p\mathrm{Fr}\nabla_\delta. \tag{15}$$

Using (15) we obtain relations between Y_i and a_{ij} :

$$\delta a_{ji} + a_{j-1,i}Y_j = a_{j,i+1}\mathrm{Fr}(Y_{i+1}). \tag{16}$$

We conclude that

$$\delta a_{ii} = 0 \tag{17}$$

$$\delta a_{i,i-1} = a_{i,i}\mathrm{Fr}Y_i - a_{i-1,i-1}Y_i \tag{18}$$

$$\delta a_{i,i-2} = a_{i,i-1}\mathrm{Fr}Y_{i-1} - a_{i-1,i-2}Y_i \tag{19}$$

$$\delta a_{30} = a_{31}\mathrm{Fr}Y_1 - a_{20}Y_3. \tag{20}$$

It follows from these formulas that $a_{i,i}$ does not depend on z and i. (Independence of z follows from the first equation; independence of i follows from second equation and from the remark that LHS of all equations vanish at $z = 0$.) Let us denote the common value of a_{ii} by ϵ; we obtain from (13) that $\epsilon = \pm 1$.

The above formulas immediately permit us to calculate every matrix element of Frobenius map up to a constant summand. To fix these constants it is sufficient to know the matrix of Frobenius map at the point $z = 0$. From (16) we can deduce that vanishing one of the elements of Frobenius matrix at $z = 0$ implies that all other elements on the same diagonal (with the same difference between indices) are zero. From the other side it is proved in [2] that in canonical coordinates q we have $a_{12} = 0$; this means that in any integral coordinate system z we have $a_{12} = 0$ at $z = 0$. (This follows from remark that $dq/dz = \pm 1$ at $z = 0$.) Applying (12),(13) we conclude that all off-diagonal entries of Frobenius matrix at $z = 0$ vanish, except, may be, a_{30}.

Similar calculations can be used to calculate the Frobenius map for $r > 1$. In this case with an appropriate choice of symplectic basis $e_0, e_1, ..., e_r, e^0, e^1, ..., e^r$ where $e_0 \in I^{3,3}, e_1, ..., e_r \in I^{2,2}, e^1, ..., e^r \in I^{1,1}, e^0 \in I^{0,0}$, Gauss-Manin connection can be represented in the following way

$$\nabla_{\delta_i} e^0 = 0, \nabla_{\delta_i} e^j =^3 Y_i^j e^0, \nabla_{\delta_i} e_j =^2 Y_{ijk} e^k, \nabla_{\delta_i} e_0 =^1 Y_i^j e_j. \tag{21}$$

We use the notation ∇_{δ_i} for the covariant derivative corresponding to $\delta_i = z_i \partial/\partial z_i$. In canonical coordinates ${}^1Y_i^j =^3 Y_i^j = \delta_i^j$, hence in arbitrary coordinate system one can express ${}^1Y_i^j =^3 Y_i^j$ in terms of mirror map. (See [4], Sec. 5.6.3 or [7].) Taking into account that in canonical coordinates ${}^2Y_i^j$ is the normalized Yukawa coupling that can expressed in terms of instanton numbers we see that all coefficients in (21) can be expressed in terms of mirror map and instanton numbers. From the other side repeating the considerations used for $r = 1$ we relate these coefficients to the Frobenius map.

Notice that we can consider also a transformation of cohomology groups over z into cohomology groups over z^q where $q = p^s$. This map denoted by $\mathrm{Fr}^{(s)}$ is induced by the s-th power of map $z \to z^p$. One can calculate this map as a composition of s Frobenius transformations. From the other side we can use the compatibility of this map with Gauss- Manin connection to calculate it in the same way as Frobenius map. Composing Frobenius transformations we obtain

$$\mathrm{Fr}^{(s)} e_i = \sum p^{(3-i)+(s-1)(3-j)} a_{ji}^{(s)} e_j \tag{22}$$

where $j \geqslant i$ and $a_{ii} = \epsilon^s$. The equation for the coefficients in (22) has the form

$$p^{s-1} a_{i+1,j}^{(s)} \mathrm{Fr}^{(s)} Y_{i+1} = \delta a_{ij}^{(s)} + a_{i,j-1}^{(s)} Y_j \tag{23}$$

3 Expression for mirror map

Integrality of mirror map was proven in [6] for quintic and in [2] in general case. In this section we will use the ideas of [2] to give an expression for mirror map in terms of Frobenius transformation. Integrality of mirror map will follow from this expression.

Our main tool will be the Artin-Hasse exponential function

$$E_p(x) = \exp(x + \frac{x^p}{p} + \frac{x^{p^2}}{p^2} + ...).$$

The expansion of this function with respect to x has integer p-adic coefficients: $E_p(x) \in \mathbb{Z}_p[[x]]$. It is easy to check that

$$\delta \log E_p(x^{pk}) - \delta \log E_p(x^k) = -kx^k = -\delta x^k.$$

Using this identity we can derive the following

Lemma 1

For every $r(x) \in \mathbb{Z}_p[[x]]$ one can find such a series $Q(x) \in x\mathbb{Z}_p[[x]]$ that $\nu(x) = \delta \log Q(x)$ obeys

$$\nu(x^p) - \nu(x) = \delta r(x).$$

To prove this lemma we give an explicit construction of $Q(x)$ in terms of Artin-Hasse exponential. Let us suppose that $r(x) = \Sigma r_k x^k$. Then it follows immediately from the properties of Artin-Hasse exponential that the power expansion of the product $Q(x) = x\Pi_k E_p(x^k)^{-r_k}$ is a series we need.

Artin-Hasse exponential can be expressed as a product:

$$E_p(x) = \Pi(1 - x^n)^{-\frac{\mu(n)}{n}}$$

where n runs over all natural numbers that are not divisible by p and μ stands for Moebius function. Using this expression we can obtain another representation of $Q(x)$:

$$Q(x) = x\Pi_k(1 - x^k)^{s_k} \tag{24}$$

where

$$s_k = \Sigma \frac{\mu(d)}{d} r_{\frac{k}{d}}$$

and d runs over all divisors of k that are not divisible by p.

In the situation considered in Sec. 2 integrality of mirror map and an explicit expression for it immediately follows from this lemma applied to the function $r(z) = \pm a_{32}(z)$. Using (18) and taking into account that $a_{ii} = \epsilon = \pm 1$ we obtain

$$\delta a_{i,i-1} = \pm(\mathrm{Fr}Y_i - Y_i).$$

Applying this relation for $i = 3$ we can say that the role of the function ν is played by the function Y_3. The mirror map q obeys $\delta \log q = Y_3$; this equation specifies q up to a constant factor. We come to a conclusion that

$$q(z) = const z \Pi_k E_p(z^k)^{\mp a_{32}^k} \tag{25}$$

where a_{32}^k denotes the coefficient of z-series $a_{32}(z)$.

One can prove that in appropriate normalization of mirror map the constant in this formula is equal to ±1. We will not discuss the calculation of this constant referring to [2].

Applying (23) we can present the expression for $q(z)$ in the form

$$q(z) = const z(1 - z^k)^{\sigma_k},$$

where

$$\sigma_k = \sum \frac{\mu(d)}{d} \epsilon a_{32}^{\frac{k}{d}}.$$

Notice that applying the above considerations to the relation (18) for arbitrary i we can prove integrality of all functions $q_i(z)$ obeying

$$\delta \log q_i = Y_i. \tag{26}$$

These functions generalizing the mirror map were introduced by Deligne [7] in geometric way; see also [4],Sec. 5.6.3. Notice that the equation (47) specifies the function $q_i(z)$ only up to a constant factor, but the geometric definition fixes this factor.

It follows from the proof of integrality of $q_i(z)$ that

$$a_{i,i-1} = \pm \log \frac{q_i(z^p)^{\frac{1}{p}}}{q_i(z)} + const \tag{27}$$

One can prove that the constant in this equation vanishes.

We have considered the mirror map only for Calabi-Yau threefolds and only in the case when $r = 1$. However, our consideration can be applied also to n-dimensional Calabi-Yau manifolds without any restriction on the dimension r of moduli space of complex structures.

In the case $r > 1$ one should apply the following multidimensional generalization of Lemma 1.

Lemma 2

Let us suppose that $r^i(z) \in \mathbb{Z}_p[[z_1, ..., z_r]]$. Then one can find such series $Q^i(z) \in z_i\mathbb{Z}_p[[z_1, ..., z_r]]$ that $\nu^i_j = \delta_j \log Q^i$ obeys

$$\mathrm{Fr}\nu^i_j - \nu^i_j = \delta_j r^i.$$

Here δ_j stands for the logarithmic derivative with repect to z_j and the Frobenius map Fr transforms $f(z_1, ..., z_r)$ into $f(z_1^p, ..., z_r^p)$.

Again we can write

$$Q^i(z) = z_i \Pi_k E_p(z^k)^{-r^i_k},$$

where r^i_k are coefficients of the power expansion of $r^i(z)$ (we consider k as multiindex).

4 Free energy and wave function

Let us consider B-model in canonical coordinate q on the moduli space $\mathcal{M}$ of complex structures. (We assume that the dimension r of $\mathcal{M}$ is equal to 1, but the generalization to the case $r > 1$ is straightforward.) Then the calculations of Sec. 2 coincide with the calculations of [1] and we obtain the following expression for Frobenius map:

$$\mathrm{Fr}e_3 = e_3, \tag{28}$$

$$\mathrm{Fr}e_2 = pe_2, \tag{29}$$

$$\mathrm{Fr}e_1 = p^2 e_1 + p^2 a_{21} e_2 + p^2 a_{31} e_3, \tag{30}$$

$$\mathrm{Fr}e_0 = p^3 e_0 + p^3 a_{20} e_2 + p^3 a_{30} e_3, \tag{31}$$

where all coefficients can be expressed in terms of $a = a_{30}$; namely

$$a_{21} = -\frac{1}{2}\delta^2 a, a_{31} = -a_{20} = \delta a.$$

Here we consider the case when all diagonal coefficients are equal to 1; in the case when the diagonal coefficients are equal to -1 we should change some signs. The symbol δ stands for logarithmic derivative with respect to q (for derivative with respect to $t = \log q$).

In canonical coordinates the function $Y_1 = Y_3$ is equal to 1 and the function $Y = Y_2$ is the Yukawa coupling (third logarithmic derivative of genus zero free energy $f = F_0$). If the Yukawa coupling has the form

$$Y(q) = c + \sum n_k q^k,$$

the genus zero free energy can be represented as

$$f(q) = ct^3 + \sum k^{-3} n_k q^k.$$

It was derived in [1] from (28-31) and compatibility of Frobenius map with Gauss-Manin connection (15) that

$$Y(q^p) - Y(q) = \frac{1}{2}\delta^3 a. \tag{32}$$

(The equations (28-31) and (32) follow from (17)-(20).) For genus zero free energy $f = F_0$ we obtain

$$p^{-3}f(q^p) - f(q) = \frac{1}{2}a. \tag{33}$$

It was shown in [1] that (32) can be used to analyze integrality of instanton numbers. It was mentioned there that the main lemma that permitted us to give p-adic interpretation of instanton numbers (=genus zero Gopakumar-Vafa invarians) can be generalized to the case of Gopakumar-Vafa invariants for arbitrary genus. Let us formulate now one of possible statements of this kind.

Recall first of all that the free energy

$$F(q) = \sum \lambda^{2g-2}F_g(q) \tag{34}$$

can be represented as a sum of the contribution of non-trivial instantons F' and the contribution of constant maps F''. The first summand can be expressed in terms of Gopakumar-Vafa invariants n_k^g:

$$F' = \sum n_k^g \frac{1}{m}(2\sin\frac{m\lambda}{2})^{2g-2}q^{mk}.$$

We sum here over all possible values of $g = 0, 1, 2, ...$ and over $k, m \in \mathbb{N}$. The second summand has the form

$$F'' = \lambda^{-2}\frac{ct^3}{6} - \frac{dt}{24}$$

where c and d are integers. Using these formulas one can represent the partition function Z as $Z'Z''$ where

$$Z' = \exp(F') = \prod (1 - \Lambda^r q^k)^{m_k^r} \tag{35}$$

$$Z'' = \exp(F''). \tag{36}$$

Here $\Lambda = e^{-i\lambda}$,

$$m_k^r = rn_k^0 + \sum_{g \geq 1+|r|} (-1)^{g-1} n_k^g \begin{pmatrix} 2g-2 \\ g-1-r \end{pmatrix}.$$

(In the formulas for free energy we disregard a constant summand, in the formulas for partition function we disregard a constant factor.)

The derivation of these formulas is based on elementary identities

$$\exp\big(\sum_m \frac{1}{m}\big(2\sin\frac{m\lambda}{2}\big)^{2g-2}q^{mk}\big) = \prod_{r=0}^{2g-2}\big(1 - \Lambda^{k-g+1}q^n\big)^{(-1)^{r+g}\binom{2g-2}{r}}$$

for $g > 0$,

$$\exp\big(\sum_m \frac{1}{m}\big(2\sin\frac{m\lambda}{2}\big)^{-2}q^{mk}\big) = \prod_{r=1}^{\infty}\big(1 - \Lambda^{-r}q^n\big)^r$$

for $g = 0$. (See [13].)

The integrality of numbers m_k^r follows from integrality of Gopakumar-Vafa invariants. Conversely, one can prove that the integrality of the numbers m_k^r together with integrality of instanton numbers n_k^0 implies the integrality of all Gopakumar-Vafa invariants n_k^g. Further, the integrality of m_k^r is equivalent to the integrality of Z' considered as an element of the ring of power series with respect to q with coefficients in Laurent series with respect to Λ. (The integrality is understood as integrality of coefficients of Laurent series.) The integrality of Z' can be expressed also in terms of p-adic reduction of Z'. (We can consider p-adic reduction of it because the coefficients in series with respect to q, Λ and, in the case, of Z'' with respect to $t = \log q, i\lambda = \log \Lambda$ are rational.) It is easy to check that for this reduction

$$\frac{Z'(\Lambda^p, q^p)}{Z'(\Lambda, q)^p} = 1 + p\zeta(\Lambda, q), \tag{37}$$

where ζ is a power series with respect to q having as coefficients Laurent series with respect to Λ ; the coefficients of these Laurent series are p-adic integers. Notice that

$$F''(p\lambda, pt) - pF''(\lambda, t) = 0, \tag{38}$$

hence

$$\frac{Z''(p\lambda, pt)}{Z''(\lambda, t)^p} = 1.$$

We see that Z also has integrality properties generalizing (37).

One should emphasize that the formula (35) is valid only for $|\Lambda| > 1$ (we used this condition in the analysis of the contribution of $g = 0$.) However, we can use the fact that Z is a function that is even with respect to λ (i.e. it does not change under the substitution $\Lambda \to \Lambda^{-1}$); therefore it can be considered as a function of $\nu = \Lambda + \Lambda^{-1} - 2 = -2\sin^2\frac{\lambda}{2}$ having a first order pole at $\nu = 0$. It is easy to derive from the above statements that the Laurent series with respect to ν that appear in Z' as coefficients in the power series with respect to q have integral coefficients. The derivation is based on the existence of one-to-one correspondence between integer polynomials with respect to ν and integer Laurent polynomials with respect to Λ that do not change under the substitution $\Lambda \to \Lambda^{-1}$. It uses also the fact that Laurent series for ν^{-1} at the point $\Lambda = \infty$ contains only negative powers of Λ appearing with integer coefficients.

Notice that the integrality properties can be expressed in terms of p-adic reduction of free energy. Namely, it follows from the above results that

$$F'(p\lambda, q^p) - pF'(\lambda, q) = p\tau, \tag{39}$$

where $\tau \in \nu^{-1}\mathbb{Z}_p[[\nu, q]]$. This statement was mentioned in [1],[2]. Using (38) we obtain that $F = F' + F''$ has the same integrality property as F'.

B-model for arbitrary genus (or, more precisely, corresponding topological string) can be obtained from genus zero B-model by means of quantization [11].

The wave function of B-model in holomorphic polarization (in polarization corresponding to the basis e_k) can be expressed in terms of free energy (34) in the following way

$$\begin{aligned}\Psi\ &= \exp\big(F(\tfrac{\lambda}{\rho}, t+\tfrac{x}{\rho}) - \lambda^{-2}(F_0(t)\rho^2 + \tfrac{\partial F_0}{\partial t}x\rho + \tfrac{1}{2}\tfrac{\partial^2 F_0}{\partial t^2}x^2)\\ &\qquad -(\tfrac{\chi}{24}-1)\log(\rho)\big). \end{aligned} \tag{40}$$

where (x, ρ) are coordinates on Lagrangian subspace spanned by e_1, e_0 and χ is an integer. (This expression was derived in [12] from the results of [10].) Let us restrict our attention to the class of wave functions having the form $\psi = e^f$ where $f = \hbar^{-1}f_0 + f_1 + \hbar f_2 + ...$ (to the class of semiclassical wave functions). In B-model $\hbar = \lambda^2$; the wave function of B-model belongs to our class. It is easy to check that a semiclassical wave function remains in the same class when we change the polarization (i.e. the group of symplectic transformation acts projectively on our space). (See [12], the end of Sec 2.) Moreover, the formulas relating the wave function $\psi = e^f$ to the wave function $\psi' = e^{f'}$ give a rational expression of f' in terms of f, hence they can be applied over any field, in particular over a field of p-adic numbers. This remark permits us to define Frobenius map on wave functions of B-model: using (28)-(31) and the formula (8) from [12] we see that we should define Frobenius map in canonical coordinates in the following way: if $\mathrm{Fr}\psi = \psi'$ then

$$f' = f + \big(\frac{1}{2\lambda^2}(a\rho^2 + \delta a\rho + \frac{1}{2}\delta^2 a x^2)\big) \tag{41}$$

As follows from (7) the Frobenius map preserves the symplectic form on cohomology up to a constant factor, hence strictly speaking it is not symplectic map; nevertheless we can use the formulas from [12] to define its action on wave functions. Notice that in (41) f' is calculated at the point q and f at the point q^p.

We conjecture that the p-adic analog of the expression Ψ obeys

$$\frac{\mathrm{Fr}\Psi}{\Psi^p} = 1 + p\zeta \tag{42}$$

where ζ is an integral expression (we understand integrality in the sense explained after the formula (37)). It follows from the above results that the integrality of Gopakumar-Vafa invariants is equivalent to this conjecture.

To verify this statement we represent Ψ as a product of three factors:

$$\Psi = \exp\big(F(\frac{\lambda}{\rho}, t+\frac{x}{\rho})\big)\exp\big(-\lambda^{-2}\big(F_0(t)\rho^2 + \frac{\partial F_0}{\partial t}x\rho + \frac{1}{2}\frac{\partial^2 F_0}{\partial t^2}x^2\big)\big)\exp\big(-(\frac{\chi}{24}-1)\log(\rho)\big).$$

It follows from our assumptions about Frobenius map and from (33) that in $\mathrm{Fr}\Psi/\Psi^p$ the contribution of the second factor in Ψ cancels with the exponent coming from (41). We obtain that (42) is equivalent to (37).

It follows from [1] that in genus zero approximation our conjecture is true; it is easy to check this fact directly.

5 Calculation of zeta-function

In Sec 2 we calculated the Frobenius map on middle-dimensional cohomology in the neighborhood of maximally unipotent boundary point. It was represented by means of a series with integral p-adic coefficients; this series converges in the open unit p-adic disk. (For simplicity we assume again that $r = 1$.) However, changing a basis in cohomology in appropriate way we can enhance the domain where our formulas make sense.

The right basis in cohomology consists of vectors $\Omega_0 = \Omega, \Omega_1 = \nabla_\delta \Omega, \Omega_2 = \nabla_\delta^2 \Omega, \Omega_3 = \nabla_\delta^3 \Omega$, where Ω stands for the cohomology class of the holomorphic 3-form. (We assume that Ω is a holomorphic function on the whole moduli space of complex structures.) We can write $\Omega = fe_0$ where $f(z)$ is a function holomorphic at $z = 0$. One can characterize $f(z)$ as a holomorphic solution of Picard-Fuchs equation. Applying (1-4) we obtain

$$\Omega_0 = fe_0, \tag{43}$$

$$\Omega_1 = \delta fe_0 + fY_1e_1, \tag{44}$$

$$\Omega_2 = \delta^2 fe_0 + 2\delta fY_1e_1 + f\delta Y_1e_1 + fY_1Y_2e_2, \tag{45}$$

$$\Omega_3 = \delta^3 fe_0 + 3\delta^2 fY_1e_1 + 3\delta f\delta Y_1e_1 + f\delta^2 Y_1e_1 + 3\delta fY_1Y_2e_2 + 2f\delta Y_1Y_2e_2 + fY_1\delta Y_2e_2 + fY_1Y_2Y_3e_3. \tag{46}$$

It is well known that zeta-function can be expressed in terms of traces on cohomology of Frobenius map and its powers at Teichmueller points. (Teichmueller points are defined as elements of the field $\mathbb{C}_p$ of complex p-adic numbers obeying $z^{p^s} = z$ for some $s \in \mathbb{N}$. In other words, these points are fixed points of powers of Frobenius map $z \to z^p$. Only at Teichmueller point the transformation $\mathrm{Fr}^{(s)}$ corresponding to a power of map $z \to z^p$ maps a cohomology group into itself and we can talk about the trace of this transformation.)

To calculate the zeta-function we should know the Frobenius map on all cohomology groups. Our results are not sufficient for this calculation if we impose only the condition of existence of non-vanishing holomorphic 3-form in the definition of Calabi-Yau threefold. (Only this condition was necessary to derive the results of Sec 2 and 3.) However, very often one includes additional conditions in the definition of Calabi-Yau manifold X; namely one requires that holomorphic forms of positive degree exist only in maximal dimension (i.e. the Hodge numbers h^{k0} vanish unless $k = 0$ or $k = \dim X$). If these conditions are included all cohomology classes on Calabi-Yau threefold, except classes in middle-dimensional cohomology, can be represented by means of algebraic cycles. If an algebraic cycle is a manifold specified by equations with rational coefficients then at a Teichmueller point an element of cohomology group dual to the cycle is an eigenvector of (a power of) Frobenius map; the eigenvalue of F^s is equal to p^{ds} where d stands for the codimension of algebraic cycle. The proof of this fact is based on the remark that one can define Frobenius map of algebraic cycles that is Poincare dual to the Frobenius map on cohomology and that the Frobenius map on algebraic cycles of the submanifold agrees with the Frobenius map on algebraic cycles of the whole manifold. If the cohomology group is spanned by algebraic cycles given by rational equations the action of Fr^s reduces to multiplication by p^{ds}. This happens, in particular, for toric manifolds and for complete intersections in toric manifolds in the case when the restriction map of two-dimensional cohomology of toric manifold into cohomology of complete intersection is surjective (for example, for quintic) .

A cohomology class that corresponds to an arbitrary algebraic subvariety still is an eigenvector with respect to Fr^s, but the eigenvalue can be equal a root of unity multiplied by p^{ds}.

This implies that the eigenvalues of the Frobenius on the second cohomology group have the form $\epsilon_i p$, where ϵ_i is a root of unity, and on the fourth cohomology group the eigenvalues are equal to $\epsilon_i^{-1}p^2$.

It is well known that zeta-function can be represented as as a quotient of two polynomials; odd-dimensional cohomology groups contribute to the numerator $P(t)$ and even-dimensional cohomology groups contribute to the denominator $Q(t)$. In the situation at hand it is easy to calculate the denominator.

If the cohomology group is spanned by algebraic cycles given by rational equations one obtains

$$Q(t) = (1-t)(1-pt)^{h^{11}}(1-p^2t)^{h^{11}}(1-p^3t). \tag{47}$$

In general case

$$Q(t) = (1-t)(1-p^3t)\prod_{1\leq i\leq h^{11}}(1-\epsilon_i pt)(1-\epsilon_i^{-1}p^2t). \tag{48}$$

All odd-dimensional cohomology groups except H^3 vanish, therefore we can obtain an expression of zeta-function in terms of Frobenius transformation on middle-dimensional cohomology calculated in Sec 2. We will not reproduce the result of this trivial, but lengthy calculation. Similar calculations can be performed in the case of elliptic curves (in this case we rederive Dwork's result [9]) and in the case of K3 surfaces (Calabi-Yau manifolds of dimension 2).

To give a flavor of formulas obtained by means of the methods described above we consider much simpler problem: how to calculate the number of points mod p^2. (We are talking about the number of points of Calabi-Yau manifold over $\mathbb{F}_p$.) As in general case we should calculate the trace of the operator Fr on the cohomology group over Teichmueller point using the basis $\Omega_0 = \Omega, \Omega_1 = \nabla_\delta\Omega, \Omega_2 = \nabla_\delta^2\Omega, \Omega_3 = \nabla_\delta^3\Omega$, but we can drop in our calculations all terms containing p^2. The calculation of this trace leads to the following result:

$$\mathrm{TrFr} = \hat{A} + \hat{B}p(1 + \frac{a}{Y_1}(\mathrm{Fr}(U) - U)) \tag{49}$$

where $a = a_{32}$ is a matrix element of Fr (see (10)), $A = fY_1^2Y_2, B = fY_1Y_2$,$U = \delta \log(f^3Y_1^2Y_2)$. We use here the notation

$$\hat{R} = \frac{\mathrm{Fr}(R)}{R}.$$

It seems that U is well defined at Teichmueller point; if this is true then $\mathrm{Fr}(U) = U$ and

$$\mathrm{TrFr} = \hat{A} + \hat{B}p.$$

Notice that one can simplify this formula using (5) . In particular, we obtain that mod p

$$\mathrm{TrFr} = \hat{A} = \widehat{f^{-1}} = \frac{f}{\mathrm{Fr}(f)}. \tag{50}$$

The above considerations permit us to say that in principle zeta-function of Calabi-Yau threefold can be expressed in terms of mirror map, instanton numbers and holomorphic period f.

There exists also an expression of zeta function in terms of all periods (solutions of Picard-Fuchs equation). Recall, that the periods $f = f_0, f_1, f_2, f_3$ can be interpreted as integrals of the form Ω over covariantly constant families of cycles or as scalar products with covariantly constant forms g_0, g_1, g_2, g_3. The scalar products $< g_a, \Omega_b >=< g_a, \nabla^b\Omega >$ can be obtained from periods by means of differentiation. We will express zeta-function in terms of matrix f_{ab} of these scalar products (period matrix). Here $0 \leq a, b \leq 3$. Let us assume that the forms g_0, g_1, g_2, g_3 constitute a symplectic basis in three-dimensional cohomology. Then $\Omega_a = r_a^b g_b$ where the matrix r is a product of symplectic matrix ε and period matrix f. The matrix, inverse to r, will be denoted by ρ. Periods are series with respect to $z, \log z$ having rational coefficients. Hence we can consider p-adic reduction of periods as well as of matrices r and ρ. (Recall that logarithm makes sense in p-adic setting.) Therefore starting with the basis Ω_i that makes sense over p-adic numbers we can construct a basis g_i of p-adic cohomology, related to Ω_i by the same formulas as in complex case. This basis also will be covariantly constant. It follows immediately from (15) that the matrix of Frobenius map in covariantly constant basis is constant; hence it is sufficient to calculate it at the point $z = 0$. The behavior of Frobenius map at this point was discussed in Sec. 2 on the base of results of [2].

We can say that the matrix of a power of Frobenius map $\mathrm{Fr}^{(s)}$ in the basis Ω_i is a product of the matrix $\mathrm{Fr}^s r$, the constant matrix of $\mathrm{Fr}^{(s)}$ at $z = 0$ and the matrix ρ. Calculating the trace of this matrix at Teichmueller points we can obtain zeta-function. Notice, that the entries of period matrix are series diverging at Teichmueller points,but the entries of the matrix of the power of Frobenius map are given by series that converge at these points.

The zeta-function of quintic was analyzed in [8]. The formula (50) agrees with [8]. It would be interesting to rederive other results of [8] using our calculation; it seems that this is possible, but not simple.

Acknowledgments

We are indebted to M. Kontsevich , A. Ogus and Daqing Wan for interesting discussions.

Both authors were partially supported by NSF grants (the first author by DMS-0505735 and the second author by DMS-0401164).

References

1. M. Kontsevich, A. Schwarz, V. Vologodsky, Integrality of instanton numbers and p-adic B-model, hep-th/0603106, Phys.Lett. B637 (2006) 97-101

2. V. Vologodsky, On integrality of instanton numbers, arXiv:0707.4617

3.A. Schwarz, I. Shapiro, Supergeometry and Arithmetic Geometry, hep-th/0605119, p-adic superspaces and Frobenius, math.NT/0605310

4.D. Cox, S. Katz, Mirror symmetry and algebraic geometry, AMS, 1999

5. D. Morisson, Mirror symmetry and rational curves on quintic threefolds: A guide for mathematicians, alg-geom/9202004; Compactifications of moduli spaces inspired by mirror symmetry,alg-geom/9304007; Mathematical aspects of mirror symmetry, alg-geom/960921

6. B. Lian, S. T. Yau, Mirror maps, modular relations and hypergeometric series, hep-th/9507151

7. P. Deligne, Local behavior of Hodge structures at infinity, in:Mirror Symmetry II (B. Greene and S.-T. Yau, eds) AMS, Providence,RI,1997, 683-699

8. P. Candelas, X. de la Ossa and F. Rodriguez-Villegas, Calabi-Yau manifolds over finite fields, I, hep-th/0012233,II,hep-th/0402133

9. B. Dwork, Normalized Period Matrices I: Plane Curves, Annals of Mathematics, 2nd Ser., Vol. 94, No. 2 (Sep., 1971) , pp. 337-388

10.Bershadsky, M., Cecotti, S., Ooguri, H., and Vafa, C., Kodaira-Spencer Theory of Gravity and Exact Results for Quantum String Amplitude, Comm. Math. Phys., 165 (1994), 211-408.

11.E. Witten , Quantum Background Independence In String Theory, hep-th/9306112.

12. A. Schwarz, Xiang Tang, Holomorphic anomaly and quantization, hep-th/0611280

13. Katz, S., Gromov-Witten, Gopakumar-Vafa, and Donaldson-Thomas invariants of Calabi-Yau threefolds, math.AG/0408266.